

Bridging the Rehabilitation Gap: A Prospective Interventional Study of Multimodal AI-based Wearable Stimulation in Subacute Stroke Patients



Sanghee Kim^{1,*} 

¹College of Nursing, Keimyung University, Daegu, Republic of Korea

Abstract:

Introduction/Objective: After discharge, many stroke patients are still in a period of active neural recovery. However, the amount of supervised rehabilitation usually decreases once they return home. This mismatch between ongoing brain plasticity and reduced therapy may affect upper limb recovery. This study examined whether a movement-responsive wearable stimulation system used at home is related to differences in upper limb motor recovery compared with usual home-based rehabilitation.

Methods: A prospective controlled study was conducted with 351 patients in the subacute stage after stroke. Participants were randomly allocated using a computer-generated sequence to either a wearable stimulation group (n = 176) or a conventional home rehabilitation group (n = 175). Upper limb motor function was assessed using the Fugl-Meyer Assessment-Upper Extremity (FMA-UE) at discharge (T0), 1 month (T1), and 3 months (T2).

Results: FMA-UE scores increased over time in both groups. However, the pattern of change differed between groups (group × time interaction, $F = 28.34$, $p < .001$). At 3 months, the mean FMA-UE score was 58.2 ± 7.5 in the wearable group and 46.1 ± 9.2 in the control group. Within the wearable group, higher device use was associated with greater motor improvement ($r = 0.72$). The between-group difference at 3 months corresponded to a large effect size (Cohen's $d = 1.4$).

Discussion: These findings suggest that movement-contingent stimulation may support motor recovery by providing feedback aligned with voluntary effort during daily activities. This approach may help sustain engagement during the post-discharge period when supervised therapy is reduced.

Conclusion: Use of a movement-responsive wearable stimulation system during home-based rehabilitation was associated with improved upper limb motor recovery during the subacute stage after stroke.

Keywords: Subacute stroke, Wearable rehabilitation, Motor function, Sensor-based feedback.

© 2026 The Author(s). Published by Bentham Open.

This is an open access article distributed under the terms of the Creative Commons Attribution 4.0 International Public License (CC-BY 4.0), a copy of which is available at: <https://creativecommons.org/licenses/by/4.0/legalcode>. This license permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

*Address correspondence to this author at the N525, College of Nursing, Keimyung University, Dalgubuldaero, Dalseo-gu, Daegu 42403, Republic of Korea; Tel: +82-53-258-7671; +82-10-9594-0128; E-mail: shkim07@gw.kmu.ac.kr

Cite as: Kim S. Bridging the Rehabilitation Gap: A Prospective Interventional Study of Multimodal AI-based Wearable Stimulation in Subacute Stroke Patients. Open Public Health J, 2026; 19: e187494452501. <http://dx.doi.org/10.2174/01187494452501260731095850>



CrossMark

Received: January 06, 2026

Revised: May 19, 2026

Accepted: June 05, 2026

Published: August 05, 2026



Send Orders for Reprints to
reprints@benthamscience.net

1. INTRODUCTION

Persistent motor impairment after stroke remains common despite improvements in acute survival rates [1-3]. Recovery patterns differ widely, particularly in relation to upper limb function. Difficulty using the affected arm interferes not only with basic daily activities but also with participation in social and occupational roles [2, 4]. For this reason, structured rehabilitation is typically initiated early in the course of care. The weeks and months following medical stabilization are often described as a period during which the nervous system remains responsive to training input [5]. Functional gains observed during this interval are not identical to those seen later in the chronic phase [6, 7]. Repetition and task-specific practice appear to influence how movement patterns reorganize [8]. Yet, the conditions under which such practice occurs frequently change once hospital-based therapy concludes.

Discharge commonly takes place before full motor recovery has been achieved [9, 10]. After returning home, therapy intensity often decreases, and patients are expected to continue exercises independently [11]. In practice, adherence varies. Some individuals maintain regular practice; others reduce activity as daily routines resume. Reduced engagement of the affected limb may contribute to compensatory strategies that persist over time. As a result, a discrepancy can emerge between recommended rehabilitation intensity and what is feasible in community settings [12, 13]. Various technological approaches have been introduced to supplement home rehabilitation [13, 14]. Wearable stimulation systems are intended to facilitate muscle activation and encourage repeated movement. Many earlier designs deliver stimulation irrespective of voluntary effort, operating in an open-loop configuration [15]. When peripheral input is not temporally aligned with motor intention, its influence on relearning processes may be limited.

Closed-loop systems attempt to address this issue by detecting movement-related signals and delivering stimulation only when active effort is identified [14, 15]. Advances in wearable sensing technologies, including inertial and electromyographic signal monitoring, have made it possible to implement such approaches outside specialized clinical environments [13]. Whether movement-contingent stimulation during the subacute stage after discharge is associated with differences in motor recovery compared with standard home instruction remains to be clarified. The present study explores this question in discharged stroke patients during the early recovery period. Closed-loop stimulation is thought to enhance motor relearning by synchronizing peripheral input with voluntary motor intention. This temporal coupling may facilitate activity-dependent plasticity and strengthen sensorimotor integration. Unlike conventional open-loop systems, the present device delivers stimulation only when movement-related signals are detected, allowing more task-relevant feedback during rehabilitation.

2. MATERIALS and METHODS

2.1. Study Design and Ethical Considerations

This investigation was carried out at three rehabilitation centers between October 2024 and October 2025. Participants were observed for three months following enrollment. Clinical evaluations were conducted by assessors who did not take part in treatment delivery. The protocol was reviewed and approved by the Institutional Review Board of the affiliated institution (IRB No. MEDR147). Written consent was obtained prior to participation. When communication was limited due to stroke-related deficits, consent procedures were completed with legally authorized representatives.

2.2. Participants

A total of 351 individuals in the recovery phase after stroke were included. Enrollment was limited to patients with a first unilateral ischemic or hemorrhagic event confirmed by imaging. At study entry, the time since onset ranged from one to three months. All participants had been discharged to home settings and demonstrated mild to moderate upper limb impairment, defined by FMA-UE scores between 15 and 55. Cognitive capacity sufficient to understand study procedures was required, reflected by Mini-Mental State Examination scores of at least 24. Patients were not enrolled if other neurological or orthopedic conditions could interfere with upper limb assessment. Additional reasons for exclusion included marked spasticity (Modified Ashworth Scale >3), implanted electronic medical devices, or skin conditions preventing electrode placement.

2.3. Allocation and Blinding

Baseline assessments were completed prior to group assignment. Allocation to the wearable intervention group ($n = 176$) or comparison group ($n = 175$) followed a computer-generated sequence prepared before recruitment began. Group assignments were revealed only after baseline measurements had been recorded. Because the intervention involved a visible device, participants were aware of their assignment. Outcome assessments, however, were conducted by occupational therapists who were not informed of group allocation.

2.4. Intervention

2.4.1. Wearable Stimulation Group

Participants assigned to the intervention received a wearable stimulation unit upon discharge. The device was positioned on the affected forearm, with electrodes placed over extensor muscle groups responsible for wrist and finger extension, including the extensor digitorum communis and extensor carpi radialis. The system incorporated motion sensing and surface electromyographic monitoring. Movement-related signals were processed continuously. When voluntary activation met predefined thresholds, electrical stimulation was triggered to assist the attempted movement. During the initial fitting, stimulation parameters were adjusted individually

to ensure comfort and effective activation. Participants were instructed to use the device for approximately one hour daily over a 12-week period. Rather than restricting use to structured exercise sessions, integration into routine activities was encouraged.

The wearable system operates in a closed-loop manner. Motion signals from the inertial sensor and surface electromyography are continuously analyzed in real time. Electrical stimulation is delivered only when voluntary movement attempts are detected and exceed predefined activation thresholds. This design allows temporal coupling between motor intention and peripheral stimulation.

Stimulation was delivered at a frequency of 30 Hz, with a pulse width of 250 μ s. The amplitude ranged from 10 to 30 mA and was adjusted individually to produce visible muscle contraction without discomfort.

2.4.2. Comparison Group

Participants in the comparison group received discharge guidance consistent with routine practice. Written materials described range-of-motion and strengthening exercises for the upper limb. These exercises were similar to those introduced during inpatient rehabilitation. Individuals were advised to continue practice at home for about 60 minutes each day for 12 weeks. No stimulation device or sensor-based feedback system was provided.

2.5. Outcome Measures

Assessments were conducted at three time points: at discharge (T0), one month (T1), and three months (T2). Upper limb motor function was evaluated using the Fugl-Meyer Assessment-Upper Extremity (FMA-UE), scored from 0 to 66. Change in score across visits was considered the primary indicator of motor recovery. Functional independence was measured with the Modified Barthel Index (MBI), which ranges from 0 to 100 and reflects performance in daily self-care activities. Health status from the participant's perspective was recorded using the EuroQol-5 Dimension (EQ-5D-5L). Adherence was tracked using different approaches. In the wearable group,

duration of use and stimulation events were stored automatically by the device. Participants in the comparison group recorded their exercise activity in written logs.

2.6. Statistical Analysis

The target sample size was estimated before recruitment began. Calculations suggested that approximately 300 participants would allow detection of modest group differences over time with 80% statistical power at a 5% alpha level. Effect sizes were calculated using Cohen's *d*. Ninety-five percent confidence intervals were also estimated. To account for possible dropout, enrollment exceeded this minimum. Analyses were conducted using SPSS software (version 28.0; IBM Corp., Armonk, NY, USA). Baseline characteristics were compared between groups using independent *t*-tests for continuous data and chi-square tests for categorical variables. Changes in outcomes across the three assessments were examined using repeated-measures models. Group was treated as the between-subject factor and time as the within-subject factor. The interaction between group and time was used to assess whether patterns of change differed across conditions. A *p*-value below .05 was considered statistically significant.

3. RESULTS

3.1. Participant Characteristics

Among 410 individuals screened at discharge, 351 fulfilled eligibility requirements and consented to participate. Fifty-nine were not enrolled because of exclusion criteria or personal decision. Group assignment resulted in 176 participants in the wearable intervention group and 175 in the comparison group. The screening and allocation process is illustrated in Fig. (1). Baseline demographic and clinical profiles were similar across groups. Age, sex distribution, BMI, affected side, cognitive function (MMSE), time since stroke onset, and initial FMA-UE scores did not differ statistically between groups (all *p* > .05). Detailed baseline information is provided in Table 1.

Table 1. Baseline characteristics of participants (N=351).

Variables	Total (N=351)	Exp. Group (n=176)	Cont. Group (n=175)	<i>p</i> -value
Age (years), Mean	68.4 $\pm$ 9.2	68.1 $\pm$ 8.9	68.7 $\pm$ 9.5	.542
Sex (male)	185 (52.7%)	93 (52.8%)	92 (52.6%)	.981
BMI (kg/m ²)	24.1 $\pm$ 3.2	24.0 $\pm$ 3.1	24.2 $\pm$ 3.3	.672
Time since Stroke (days)	52.3 $\pm$ 14.8	51.6 $\pm$ 14.2	53.0 $\pm$ 15.3	.210
Stroke Type	-	-	-	.844
Ischemic	247 (70.4%)	123 (69.9%)	124 (70.9%)	-
Hemorrhagic	104 (29.6%)	53 (30.1%)	51 (29.1%)	-
Affected side (Right)	182 (51.9%)	91 (51.7%)	91 (52.0%)	.954
MMSE score	26.8 $\pm$ 1.9	26.7 $\pm$ 1.8	26.9 $\pm$ 2.0	.441
Baseline FMA-UE Score	32.5 $\pm$ 8.4	32.1 $\pm$ 8.2	32.9 $\pm$ 8.6	.385

Note: Values are presented as mean $\pm$ standard deviation (SD) or number (%). BMI: body mass index; MMSE: Mini-Mental State Examination; FMA-UE: Fugl-Meyer Assessment-Upper Extremity.




Fig. (1). Participant flow from screening to follow-up.

3.2. Motor Performance and Functional Outcomes

Upper limb motor scores changed across follow-up in both groups. Patterns of change, however, were not identical. Repeated-measures analysis identified a significant interaction between group and time ($F = 28.34$, $p < .001$). Between one and three months, improvement appeared to slow in the comparison group, while the intervention group continued to show progressive gains. At the three-month assessment, mean FMA-UE values were 58.2 ± 7.5 in the intervention group and 46.1 ± 9.2 in the comparison group. The difference between groups at this point reached statistical significance ($p < .001$). Detailed outcome data are presented in Table 2. Changes

in FMA-UE scores over time are shown in Fig. (2A), whereas changes in MBI scores are presented in Fig. (2B). Activities of daily living demonstrated a similar directional pattern. MBI scores increased during follow-up, with a significant group-by-time interaction ($F = 15.67$, $p < .001$). At three months, the intervention group recorded a mean score of 92.1 compared with 79.5 in the comparison group. Quality-of-life scores also improved over time. Between-group differences were smaller in magnitude than those observed for motor performance but reached statistical significance at follow-up ($p < .05$). The mean difference in FMA-UE at 3 months was 12.1 points (95% CI: 10.2-14.0). All outcome values are summarized in Table 2.

Table 2. Changes in upper limb motor function and quality of life over time.

Variable	Group	Baseline (T0)	1 Month (T1)	3 Months (T2)	Mean Diff (T2)	95% CI	Effect Size	F (Interaction)
FMA-UE	Exp.	32.1 ± 8.2	45.3 ± 9.1	58.2 ± 7.5	12.1**	10.2-14.0	d=1.40	28.34*
(Motor)	Cont.	32.9 ± 8.6	38.5 ± 8.8	46.1 ± 9.2	-	-	-	-
MBI	Exp.	65.2 ± 10.5	78.4 ± 11.2	92.1 ± 8.4	8.12*	9.8-15.4	d=1.20	15.67*
(ADL)	Cont.	64.8 ± 10.1	70.1 ± 10.8	79.5 ± 11.5	-	-	-	-

Note: Exp = intervention group; Cont = comparison group; CI: confidence interval. Values are presented as mean $\pm$ standard deviation (SD). Mean Diff (T2): between-group difference at 3 months (Exp - Cont). Effect size calculated using Cohen's d. * $p < .05$; ** $p < .01$; *** $p < .001$.




Fig. (2). Changes in motor function and functional independence.

Note: **Figure 2A** presents FMA-UE scores measured at discharge (T0), 1 month (T1), and 3 months (T2). **Figure 2B** presents Modified Barthel Index (MBI) scores measured at the same assessment points. Values are expressed as mean $\pm$ standard deviation (SD). Dashed gridlines are shown for visual reference. *** $p < .001$ for the between-group comparison at 3 months.




Fig. (3). Association between device use and motor recovery.

3.3. Association Between Device Use and Motor Recovery

Within the wearable group, variation in recorded device use was examined in relation to motor change over three months. Compliance reflected the proportion of activation events triggered during valid movement attempts. Motor improvement was expressed as the difference in FMA-UE score between discharge and the three-month assessment. A positive association was observed between recorded device use and change in FMA-UE score ($r = 0.72, p < .001$). Individuals with higher activation rates generally showed larger increases in motor performance. The distribution of this relationship is presented in Fig. (3).

4. DISCUSSION

The weeks following discharge often introduce a shift in how rehabilitation is experienced. During inpatient care, repetition is structured, supervised, and externally regulated. Once at home, the same level of intensity is difficult to maintain. Neurological recovery, however, does not pause at discharge [16]. The early post-stroke period continues to be characterized by adaptive processes that may respond to practice input [17]. The contrast between ongoing biological responsiveness and reduced therapeutic structure raises practical questions about how recovery environments shape functional change.

Home exercise programs depend heavily on self-management. Motivation, daily routine, fatigue, and family support all influence how frequently prescribed exercises are performed. Prior work has noted that adherence may

decline when repetition lacks variation or external reinforcement [18]. In this context, movement-contingent assistance introduces a different form of engagement. Rather than providing stimulation continuously, the system responds only when voluntary effort is detected. This conditional activation changes the temporal pattern of feedback and may influence how movements are repeated and perceived [19].

The association between recorded device use and motor change adds nuance to this interpretation. Rehabilitation outcomes are not determined solely by duration of practice [20]. The quality of repetition—its timing, attentional demands, and contextual integration—also plays a role. When assistance occurs during an intentional movement attempt, repetition may be more closely linked to goal-directed activity. Although direct neural mechanisms were not examined, the observed pattern aligns with conceptual models emphasizing contingent feedback in motor relearning [21]. One possible explanation is that movement-contingent stimulation enhances sensorimotor coupling by reinforcing motor cortex activation during voluntary effort. This timing-dependent feedback may support activity-dependent plasticity and improve motor relearning.

It is also important to consider that behavioral factors can influence engagement. The presence of monitoring may alter routine behavior. Participants aware of device recording may interact differently with the intervention. In addition, the social environment at home varies widely. Caregiver involvement, encouragement, and logistical support may shape daily activity patterns. Such contextual

differences are rarely uniform across participants and may contribute to variability in outcome trajectories [22].

The broader context of stroke rehabilitation further frames these findings. Discontinuity between hospital-based therapy and community follow-up has been described as a persistent structural challenge. Access to outpatient or community rehabilitation services varies across regions, and extended inpatient stays are often limited by resource allocation [23]. Evidence suggests that home-based rehabilitation can be as effective as center-based programs when adequate intensity and guidance are maintained [24]. In this context, technology-supported approaches may represent one method of sustaining structured engagement when direct supervision decreases. Their role, however, should be viewed as complementary rather than substitutive, given the multidimensional needs of stroke survivors.

Interpretation remains cautious. Observational associations cannot establish causal pathways. Variability in adherence measurement methods between groups and differences in personal circumstances may have influenced the results. Follow-up was confined to the early recovery window, and trajectories beyond this period remain uncertain. Longer observation would clarify whether early differences narrow, persist, or evolve as recovery stabilizes.

5. LIMITATIONS

Several aspects of the study design should be considered. Participants knew which group they were assigned to, and this awareness may have influenced how they engaged with the program. In behavioral rehabilitation research, separating intervention effects from motivational factors is not straightforward. Adherence was captured differently in each group. Device activity was recorded automatically in the intervention group, whereas participants in the comparison group documented their exercises in written logs. These approaches are not directly equivalent and may have affected how engagement was represented. Follow-up covered only the first three months after discharge. Changes observed during this stage may not reflect longer-term patterns. In addition, the analysis focused on functional scores and recorded device activity. Factors such as ease of use in home environments, caregiver support, and the day-to-day demands of managing equipment were not examined in detail. These practical considerations may influence how similar systems perform outside controlled study settings.

CONCLUSION

The use of a movement-responsive wearable system during the early months after discharge was associated with differences in motor and functional outcomes compared with standard home guidance. Structured, effort-contingent feedback may help sustain engagement during a period when supervised therapy decreases. How such approaches perform over longer periods and across varied home contexts remains to be clarified.

AUTHORS' CONTRIBUTIONS

The authors confirm contribution to the paper as follows: K.S.H.: Conceived and designed the study, conducted data collection and analysis, interpreted the findings, and prepared the manuscript. The author reviewed and approved the final version for submission.

LIST OF ABBREVIATIONS

ADL	= Activities of Daily Living
ECR	= Extensor Carpi Radialis
EDC	= Extensor Digitorum Communis
EQ-5D	= EuroQol-5 Dimension
FES	= Functional Electrical Stimulation
FMA-UE	= Fugl-Meyer Assessment-Upper Extremity
IMU	= Inertial Measurement Unit
MBI	= Modified Barthel Index
sEMG	= Surface Electromyography

ETHICS APPROVAL AND CONSENT TO PARTICIPATE

Ethical approval was obtained from the Institutional Review Board of the affiliated institution (Approval Date: October 2024).

HUMAN AND ANIMAL RIGHTS

All human research procedures followed were in accordance with the ethical standards of the committee responsible for human experimentation (institutional and national), and with the Helsinki Declaration of 1975, as revised in 2013.

CONSENT FOR PUBLICATION

Participation was voluntary, and written consent was secured prior to study procedures. When communication limitations were present due to stroke-related impairments, consent was provided by legally authorized representatives. Data were coded prior to analysis, and identifying information was removed to protect participant confidentiality.

STANDARDS OF REPORTING

STROBE guidelines were followed.

AVAILABILITY OF DATA AND MATERIALS

All the data and supporting material is available within the article.

FUNDING

None.

CONFLICT OF INTEREST

The author declares no conflict of interest, financial or otherwise.

ACKNOWLEDGEMENTS

Declared none.

REFERENCES

- [1] Tang Q, Yang X, Sun M, *et al.* Research trends and hotspots of post-stroke upper limb dysfunction: a bibliometric and visualization analysis. *Front neurol* 2024; 15: 1449729. <http://dx.doi.org/10.3389/fneur.2024.1449729> PMID: 39416663
- [2] Ingram LA, Butler AA, Brodie MA, Lord SR, Gandevia SC. Quantifying upper limb motor impairment in chronic stroke: a physiological profiling approach. *J appl physiol* (Bethesda Md, : 1985) 2021; 131(3): 949-65. <http://dx.doi.org/10.1152/japplphysiol.00078.2021> PMID: 34264125
- [3] Chen X, Catherine Chan KL, Wang X, *et al.* Simplified assessment of upper limb dysfunction after stroke: decision tree analysis based on the International Classification of Functioning, Disability and Health. *Eur j phys rehabil med* 2025; 61(5): 735-43. <http://dx.doi.org/10.23736/S1973-9087.25.08976-2> PMID: 41324392
- [4] Zhou X, Du M, Dai X, Zhu S, Zhou L, Li X. Intervention patterns and preliminary effectiveness on Social Participation following stroke: a scoping review. *BMC neurol* 2023; 23(1): 275. <http://dx.doi.org/10.1186/s12883-023-03250-2> PMID: 37464300
- [5] Li X, He Y, Wang D, Rezaei MJ. Stroke rehabilitation: from diagnosis to therapy. *Front neurol* 2024; 15: 1402729. <http://dx.doi.org/10.3389/fneur.2024.1402729> PMID: 39193145
- [6] Poomalai G, Prabhakar S, Sirala Jagadesh N. Functional Ability and Health Problems of Stroke Survivors: An Explorative Study. *Cureus* 2023; 15(1): e33375. <http://dx.doi.org/10.7759/cureus.33375> PMID: 36751244
- [7] Altuntaş O, Taş S, Çetin A. An investigation of the factors that influence functional improvement in stroke rehabilitation. *Turk j med sci* 2021; 51(3): 1448-54. <http://dx.doi.org/10.3906/sag-2101-94> PMID: 33705646
- [8] Chen G, Wu M, Chen J, *et al.* Biomarkers associated with functional improvement after stroke rehabilitation: a systematic review and meta-analysis of randomized controlled trials. *Front neurol* 2023; 14: 1241521. <http://dx.doi.org/10.3389/fneur.2023.1241521> PMID: 37731856
- [9] Ruksakulpiwat S, Benjasirisan C, Phianhasin L, *et al.* Effectiveness of Discharge Planning Interventions for Stroke and Heart Conditions: A Systematic Review of Interventional Studies. *J Multidiscip Healthc* 2025; Volume 18: 7521-37. <http://dx.doi.org/10.2147/jmdh.s563476>
- [10] , Rahmi U, Fitriana LA, *et al.* The Impact of Discharge Planning on Enhancing Independence in Ischaemic Stroke Patients: A Post-hospitalisation Rehabilitation Approach. *Malays J Med Sci* 2025; 32(2): 99-107. <http://dx.doi.org/10.21315/mjms-01-2025-027>
- [11] Chin LF, Rosbergen ICM, Hayward KS, Brauer SG. A self-directed upper limb program during early post-stroke rehabilitation: A qualitative study of the perspective of nurses, therapists and stroke survivors. *PloS one* 2022; 17(2): e0263413. <http://dx.doi.org/10.1371/journal.pone.0263413> PMID: 35120167
- [12] Hung SH, Ackerley S, Connell LA, *et al.* Real-World Experiences of Therapy Staff Implementing an Intensive Rehabilitation Protocol in Canadian Stroke Inpatient Rehabilitation Settings: A Multi-Site Survey Study. *Phys ther* 2025; 105(10) <http://dx.doi.org/10.1093/ptj/pzaf111> PMID: 41139309
- [13] Sweeney G, Boyd F, Keogh M, *et al.* A technology-enriched approach to increasing rehabilitation dose after stroke: Clinical feasibility study. *Clin rehabil* 2025; 39(6): 740-9. <http://dx.doi.org/10.1177/02692155251333542> PMID: 40247700
- [14] Hong R, Li B, Bao Y, Liu L, Jin L. Therapeutic robots for post-stroke rehabilitation. *Med rev* (2021) 2024; 4(1): 55-67. <http://dx.doi.org/10.1515/mr-2023-0054> PMID: 38515779
- [15] Fong KNK, Pak JPY, Koo AHL, *et al.* Closed-loop versus open-loop "remind-to-move" treatment using wearables for hemiparetic upper extremity in patients after stroke: A proof-of-concept study. *Wearable technol* 2025; 6: e33. <http://dx.doi.org/10.1017/wtc.2025.10017> PMID: 40692810
- [16] Lee J, Kim G. Functional Recovery in Acute and Subacute Stroke Patients With or Without Post-stroke Fatigue. *Brain NeuroRehabilit* 2024; 17(3): e22. <http://dx.doi.org/10.12786/bn.2024.17.e22> PMID: 39649708
- [17] Du M, Chen L, Li Y, *et al.* Effect of exercise-based interventions on stroke rehabilitation: an umbrella review of systematic reviews and meta-analyses. *J neuroeng rehabil* 2025; 22(1): 240. <http://dx.doi.org/10.1186/s12984-025-01781-y> PMID: 41250143
- [18] Nascimento LR, Rocha RJ, Boening A, Ferreira GP, Perovano MC. Home-based exercises are as effective as equivalent doses of centre-based exercises for improving walking speed and balance after stroke: a systematic review. *J physiother* 2022; 68(3): 174-81. <http://dx.doi.org/10.1016/j.jphys.2022.05.018> PMID: 35753966
- [19] Langerak AJ, Regterschot GRH, Evers M, *et al.* A Sensor-Based Feedback Device Stimulating Daily Life Upper Extremity Activity in Stroke Patients: A Feasibility Study. *Sens (Basel Switz)* 2023; 23(13) <http://dx.doi.org/10.3390/s23135868> PMID: 37447718
- [20] Kwakkel G, Stinear C, Essers B, *et al.* Motor rehabilitation after stroke: European Stroke Organisation (ESO) consensus-based definition and guiding framework. *Eur Stroke J* 2023; 8(4): 880-94. <http://dx.doi.org/10.1177/23969873231191304>
- [21] Sarzyńska-Długosz I. An optimal model of long-term post-stroke care. *Front Neurol* 2023; 14 <http://dx.doi.org/10.3389/fneur.2023.1129516>
- [22] Platz T. Evidence-Based Guidelines and Clinical Pathways in Stroke Rehabilitation-An International Perspective. *Front neurol* 2019; 10: 200. <http://dx.doi.org/10.3389/fneur.2019.00200> PMID: 30930832
- [23] Nascimento LR, Gaviorno LF, de Souza Brunelli M, Gonçalves JV, Arêas FZDS. Home-based is as effective as centre-based rehabilitation for improving upper limb motor recovery and activity limitations after stroke: A systematic review with meta-analysis. *Clin Rehabil* 2022; 36(12): 1565-77. <http://dx.doi.org/10.1177/02692155221121015>
- [24] Basheikh MA, Badahdah AA. Efficacy of Home-Based Physical Exercise in Stroke Survivors: A Systematic Review and Meta-Analysis of Randomized Controlled Trials. *Arch rehabil res clin transl* 2025; 7(4): 100494. <http://dx.doi.org/10.1016/j.arrct.2025.100494> PMID: 41477087

DISCLAIMER: The above article has been published, as is, ahead-of-print, to provide early visibility but is not the final version. Major publication processes like copyediting, proofing, typesetting and further review are still to be done and may lead to changes in the final published version, if it is eventually published. All legal disclaimers that apply to the final published article also apply to this ahead-of-print version.